

| Tree # | Common Name | Botanical Name | DBH (in) | Condition (%) | Height (ft) | Width (ft) | Clear Trunk (ft) | Canopy Coverage (sq ft) | Notes |
|--------|---------------|---------------------------|----------|---------------|-------------|------------|------------------|-------------------------|---|
| 1 | Bismarck Palm | <i>Bismarckia nobilis</i> | | 90% | 25 | 14 | 15 | 154 | |
| 2 | Royal Palm | <i>Roystonea regia</i> | | 50% | 28 | 12 | 20 | 113 | Severe penciling in upper portion of trunk. Chlorotic crown. Trunk has oozing cracks. |
| 3 | Bismarck Palm | <i>Bismarckia nobilis</i> | | 90% | 29 | 16 | 17 | 201 | |
| 4 | Royal Palm | <i>Roystonea regia</i> | | 60% | 35 | 14 | 28 | 154 | Vertical cracks along trunk. Mild chlorosis in crown. |
| 5 | Bismarck Palm | <i>Bismarckia nobilis</i> | | 90% | 22 | 16 | 12 | 201 | |

AAC

PZ24-12000027
06/02/2026

AAC

PZ24- 12000027
09/10/2025

| | | | | | | | | | |
|----|-------------------|------------------------------|----|-----|----|----|----|-----|---|
| 6 | Sabal Palm | <i>Sabal palmetto</i> | | 80% | 12 | 12 | 4 | 113 | Limited root zone. Utility pole conflict. |
| 7 | Sabal Palm | <i>Sabal palmetto</i> | | 70% | 9 | 10 | 11 | 79 | Limited root zone. Tip dieback/chlorosis on most fronds. |
| 8 | Silver Buttonwood | <i>Conocarpus erectus</i> | 10 | 55% | 16 | 8 | | 50 | Major codominance in lower half. Low live crown ratio. Limited root zone. |
| 9 | Silver Buttonwood | <i>Conocarpus erectus</i> | 10 | 75% | 15 | 10 | | 79 | Poor structure. Limited root zone. Good health and vigor. |
| 10 | False Tamarind | <i>Lysiloma latisiliquum</i> | 20 | 55% | 14 | 20 | | 314 | Poor structure. Multiple codominant trunks in lower half with included bark. Limited root zone in all directions. Roots lifting pavement and curb. Poor vigor. Dieback of limbs 1-3" in diameter throughout interior. |
| 11 | Silver Buttonwood | <i>Conocarpus erectus</i> | 10 | 80% | 16 | 14 | | 154 | Limited root zone. Over-extended branches. |

| | | | | | | | | | |
|----|-------------------|-------------------------------|----|-----|----|----|---|-----|--|
| 12 | Silver Buttonwood | <i>Conocarpus erectus</i> | 10 | 60% | 18 | 14 | | 154 | Poor structure. Multiple codominant trunks in lower half. Sparse foliage within crown. Overextended branches. |
| 13 | Montgomery Palm | <i>Veitchia montgomeryana</i> | | 80% | 10 | 5 | 7 | 20 | |
| 14 | Silver Buttonwood | <i>Conocarpus erectus</i> | 13 | 80% | 20 | 18 | | 254 | Codominant in lower half. Slight lean to the North. Good health and vigor. |
| 15 | Silver Buttonwood | <i>Conocarpus erectus</i> | 9 | 75% | 12 | 14 | | 154 | Limited root zone. Overhead wire conflict. Vertically-oriented/weakly-attached branches within crown. Good health and vigor. |
| 16 | Gumbo Limbo | <i>Bursera simaruba</i> | 6 | 60% | 15 | 10 | | 79 | Overhead wire conflict. Poor structure. Limited root zone. |
| 17 | Silver Buttonwood | <i>Conocarpus erectus</i> | 6 | 60% | 11 | 16 | | 201 | Overhead wire conflict. Poor structure. Limited root zone. |

| | | | | | | | | | |
|----|----------------|----------------------------------|---|-----|----|----|----|----|-----------------------------------|
| 18 | Carrotwood | <i>Cupaniopsis anacardioides</i> | 3 | 20% | 10 | 6 | | 28 | FISC category I invasive species. |
| 19 | Umbrella Tree | <i>Schefflera actinophylla</i> | 5 | 50% | 10 | 6 | | 28 | FISC category I invasive species. |
| 20 | Areca Palm | <i>Dypsis lutescens</i> | | 80% | 10 | 10 | 6 | 79 | Multi-trunk cluster. |
| 21 | Christmas Palm | <i>Adonidia merrillii</i> | | 80% | 14 | 4 | 10 | 13 | |
| 22 | Christmas Palm | <i>Adonidia merrillii</i> | | 80% | 18 | 4 | 14 | 13 | |
| 23 | Christmas Palm | <i>Adonidia merrillii</i> | | 80% | 14 | 10 | 10 | 79 | Double. |

| | | | | | | | | | |
|----|----------------|---------------------------|--|-----|----|----|----|----|---------|
| 24 | Christmas Palm | <i>Adonidia merrillii</i> | | 80% | 18 | 4 | 14 | 13 | |
| 25 | Christmas Palm | <i>Adonidia merrillii</i> | | 80% | 18 | 4 | 14 | 13 | |
| 26 | Christmas Palm | <i>Adonidia merrillii</i> | | 80% | 12 | 4 | 8 | 13 | |
| 27 | Christmas Palm | <i>Adonidia merrillii</i> | | 80% | 18 | 4 | 14 | 13 | |
| 28 | Christmas Palm | <i>Adonidia merrillii</i> | | 80% | 16 | 10 | 12 | 79 | Double. |
| 29 | Christmas Palm | <i>Adonidia merrillii</i> | | 80% | 14 | 4 | 10 | 13 | |

| | | | | | | | | | |
|----|-------------------|--------------------------------|----|-----|----|----|----|-----|-----------------------------------|
| 30 | Umbrella Tree | <i>Schefflera actinophylla</i> | | 50% | 16 | 8 | | 50 | FISC category I invasive species. |
| 31 | Christmas Palm | <i>Adonidia merrillii</i> | | 80% | 20 | 4 | 16 | 13 | |
| 32 | Christmas Palm | <i>Adonidia merrillii</i> | | 80% | 18 | 10 | 14 | 79 | Double. |
| 33 | Christmas Palm | <i>Adonidia merrillii</i> | | 80% | 16 | 4 | 12 | 13 | |
| 34 | Silver Buttonwood | <i>Conocarpus erectus</i> | 12 | 80% | 10 | 12 | | 113 | Limited root zone. |
| 35 | Christmas Palm | <i>Adonidia merrillii</i> | | 80% | 18 | 10 | 14 | 79 | Double |

AAC

PZ24-12000027
06/02/2026

AAC

PZ24- 12000027
09/10/2025

| | | | | | | | | | |
|----|-----------------------|-------------------------------|----|-----|----|----|---|-----|---|
| 36 | Christmas Palm | <i>Adonidia merrillii</i> | | 80% | 10 | 4 | 6 | 13 | |
| 37 | False Tamarind | <i>Lysiloma latisiliquum</i> | 13 | 70% | 18 | 16 | | 201 | Limited root zone. Codominant in lower half. |
| 38 | Christmas Palm | <i>Adonidia merrillii</i> | | 60% | 12 | 5 | 7 | 20 | Single. Canopy competition with Brazilian Pepper. |
| 39 | Brazilian Beauty Leaf | <i>Calophyllum inophyllum</i> | 12 | 80% | 20 | 20 | | 314 | Limited root zone. |
| 40 | Brazilian Beauty Leaf | <i>Calophyllum inophyllum</i> | 11 | 80% | 15 | 20 | | 314 | Limited root zone. |
| 41 | Live Oak | <i>Quercus virginiana</i> | 6 | 75% | 16 | 14 | | 154 | Limited root zone. |

| | | | | | | | | | |
|----|-----------------------|-------------------------------|----|-----|----|----|----|-----|---|
| 42 | Brazilian Beauty Leaf | <i>Calophyllum inophyllum</i> | 10 | 75% | 20 | 20 | | 314 | Limited root zone. Mildly chlorotic foliage. |
| 43 | Live Oak | <i>Quercus virginiana</i> | 5 | 60% | 18 | 14 | | 154 | Limited root zone. Poor vigor. |
| 44 | Brazilian Beauty Leaf | <i>Calophyllum inophyllum</i> | 12 | 60% | 20 | 15 | | 177 | Limited root zone. Girdling roots. Tip dieback. |
| 45 | Brazilian Beauty Leaf | <i>Calophyllum inophyllum</i> | 12 | 70% | 25 | 15 | | 177 | Limited root zone. Plunging and girdling roots. |
| 46 | Coconut Palm | <i>Cocos nucifera</i> | | 75% | 40 | 16 | 34 | 201 | Minor mechanical trunk damage. |
| 47 | Sabal Palm | <i>Sabal palmetto</i> | | 80% | 12 | 14 | 5 | 154 | |

| | | | | | | | | | |
|----|-----------------------|-------------------------------|----|-----|----|----|--|-----|--|
| 48 | Brazilian Beauty Leaf | <i>Calophyllum inophyllum</i> | 8 | 10% | 14 | 10 | | 79 | Severe decline and dieback. |
| 49 | Brazilian Beauty Leaf | <i>Calophyllum inophyllum</i> | 12 | 70% | 12 | 18 | | 254 | |
| 50 | Brazilian Beauty Leaf | <i>Calophyllum inophyllum</i> | 11 | 65% | 16 | 16 | | 201 | Tip dieback. |
| 51 | Brazilian Beauty Leaf | <i>Calophyllum inophyllum</i> | 11 | 65% | 20 | 20 | | 314 | Tip dieback. |
| 52 | False Tamarind | <i>Lysiloma latisiliquum</i> | 12 | 55% | 18 | 14 | | 154 | Severely limited root zone. Dieback of branches 1-3" in diameter. Hanging branches. Cracking along main trunk. |
| 53 | Live Oak | <i>Quercus virginiana</i> | 8 | 60% | 18 | 16 | | 201 | Limited root zone. Girdling roots. |

| | | | | | | | | | |
|----|------------------|--------------------------------|----|-----|----|----|--|-----|---|
| 54 | Live Oak | <i>Quercus virginiana</i> | 10 | 70% | 18 | 18 | | 254 | Limited root zone. Tip dieback. |
| 55 | Live Oak | <i>Quercus virginiana</i> | 10 | 50% | 18 | 20 | | 314 | Limited root zone. Dieback. Poor vigor. |
| 56 | Live Oak | <i>Quercus virginiana</i> | 11 | 60% | 20 | 20 | | 314 | Limited root zone. Lifting roots. Dieback of interior foliage within crown. |
| 57 | Live Oak | <i>Quercus virginiana</i> | 8 | 50% | 18 | 15 | | 177 | Limited root zone. Girdling roots. Tip dieback. Poor vigor. |
| 58 | Live Oak | <i>Quercus virginiana</i> | 11 | 70% | 20 | 20 | | 314 | Limited root zone. Overextended branches. |
| 59 | Yellow Poinciana | <i>Peltophorum pterocarpum</i> | 20 | 55% | 18 | 20 | | 314 | Poor structure. Codominance in lower half with included bark. Severely limited root zone. Lifting roots. Visual evidence of internal decay. |

| | | | | | | | | | |
|----|-----------------|-----------------------------|----|-----|----|----|--|-----|--|
| 60 | Live Oak | <i>Quercus virginiana</i> | 10 | 80% | 15 | 15 | | 177 | Limited root zone. Over-extended branches. |
| 61 | Live Oak | <i>Quercus virginiana</i> | 10 | 70% | 14 | 16 | | 201 | Limited root zone. Tip dieback. |
| 62 | Live Oak | <i>Quercus virginiana</i> | 10 | 60% | 16 | 16 | | 201 | Limited root zone. Tip dieback. Circling roots. |
| 63 | Simpson Stopper | <i>Myrcianthes fragrans</i> | 12 | 80% | 12 | 10 | | 79 | Multi-trunk. Limited root zone. Canopy and root competition. |
| 64 | Simpson Stopper | <i>Myrcianthes fragrans</i> | 8 | 80% | 10 | 6 | | 28 | Multi-trunk. Hedge-sheared. |
| 65 | Live Oak | <i>Quercus virginiana</i> | 10 | 30% | 14 | 16 | | 201 | Very poor structure. No central leader. Weak attachments. |

| | | | | | | | | | |
|----|-----|-----|----|----|----|----|--|-----|-------|
| 66 | N/A | N/A | 12 | 0% | 20 | 15 | | 177 | Dead. |
|----|-----|-----|----|----|----|----|--|-----|-------|